



Well-Posedness of Jump Problem on Non-rectifiable Curves

David B. Katz² · Boris A. Kats¹

Received: 25 September 2017 / Accepted: 8 November 2018
© Springer Nature Switzerland AG 2018

Abstract

We investigate the well-posedness of a jump problem on non-rectifiable curves with Hölder boundary data. The jump problem is a special case of Riemann boundary value problem. However, the classic results are obtained in the case of piecewise smooth boundary. There are some results for non-rectifiable boundary (see our previous papers in bibliography), but still there is a large field of questions waiting for answers.

Keywords Well-posedness · Jump problem · Non-rectifiable curve

Introduction

We consider the following boundary value problem for analytic functions. We have a simple closed curve Γ on the complex plane, which divides it into domains D^+ and D^- , $\infty \in D^-$. We need to find all analytic in D^+ and D^- functions $\Phi^\pm(z)$ such that difference of their boundary values equals to the given function f . It is a well known special case of the Riemann boundary value problem, see [1–3]. A great body of publications are dealing with applications of these problems in mathematical physics, theory of integral equations and other fields.

In the monographs mentioned above the curve Γ is assumed smooth or piecewise smooth, and the jump f is Hölder-continuous on Γ . In this case the Cauchy type

Communicated by Laurent Baratchart.

The research is partially supported by Russian Foundation for Basic Researches and Government of Republic Tatarstan, Grant No. 18-41-160003.

✉ Boris A. Kats
katsboris877@gmail.com

David B. Katz
kazdavid89@gmail.com

¹ Kazan (Volga Region) Federal University, Kremlevskaya str. 18, Kazan, Russia 420138

² Mathematical Modeling of Nonequilibrium Processes in Oil and Gas Production Laboratory, Kazan (Volga Region) Federal University, Kremlevskaya str. 18, Kazan, Russia 420138